

**Blake M. Bodendorfer, MD**  
[omahasportsdoc.com](http://omahasportsdoc.com)

## **Rehabilitation Protocol Nonoperative Radial Head Fracture**

### **Phase I: Immobilization (3-7 days)**

Sling should be worn for comfort for only 3-7 days.

- Grip and wrist/hand AROM immediately.
- Begin putty/grip/ 6 pack exercises

### **Phase II: Early ROM (1 to 2 weeks)**

- Remove sling at all times
- Initiate elbow exercise program 5 times per day:
- Passive and active elbow ROM as tolerated:
  - o flexion/extension/pronation/supination
- ROM by end of 2nd week should be at least 15 to 115 degrees
- Begin putty/grip exercises
- Begin isometric strengthening exercises for the elbow and wrist

### **Phase III: Increase ROM and function (2 to 6 weeks)**

- Continue elbow active and active assisted ROM exercises.
- Full flexion and extension should be achieved by the end of 6 weeks.
- Begin light isotonic strengthening of flexion and extension
- Assess and maintain shoulder and wrist strength and ROM

### **Phase IV: Full ROM and function (>6 weeks)**

- Full pronation and supination should be achieved by the end of the 8th week.
- Progressively increase isotonic strengthening in flexion/extension and pronation/supination.
- Work on any deficits